

Table B.8: Top 10 leading causes of reported hospital deaths* by ICD10 4-character code and gender, 2017

| Top 10 leading causes of death | | | | | |
|---|--------|-----|---|--------|-----|
| Males | | | Females | | |
| Description | ICD-10 | % | Description | ICD-10 | % |
| Malignant neoplasm of bronchus or lung | C34.9 | 4.6 | Malignant neoplasm of breast | C50.9 | 2.7 |
| Acute myocardial infarction | I21.9 | 3.0 | Malignant neoplasm of bronchus or lung | C34.3 | 2.2 |
| Pneumonia | J18.9 | 2.2 | Acute myocardial infarction | I21.9 | 2.0 |
| Exposure to unspecified factor, unspecified place | X59.9 | 2.1 | Pneumonia | J18.9 | 1.7 |
| Acute renal failure | N17.9 | 1.4 | Acute renal failure | N17.9 | 1.5 |
| Intracerebral haemorrhage | I61.9 | 1.2 | Malignant neoplasm of colon | C18.9 | 0.9 |
| Malignant neoplasm of colon | C18.9 | 1.2 | Malignant neoplasm of pancreas | C25.9 | 0.8 |
| Malignant neoplasm of pancreas | C25.9 | 1.1 | Intracerebral haemorrhage | I61.9 | 0.8 |
| Malignant neoplasm of prostate | C61 | 1.1 | Stroke | I64 | 0.7 |
| Malignant neoplasm of bladder | C67.9 | 0.8 | Exposure to unspecified factor, unspecified place | X59.9 | 0.7 |

Source: Hospital Mortality System (HMS), Statistics Department, MOPH (on 12-11-2018)

*The underlying cause of death was selected using the IRIS software, a coding system based on international standard